

Horizon 2020 Programme

INFRAIA-02-2017 Integrating Activities for Starting Communities



SmartCow: an integrated infrastructure for increased research capability and innovation in the European cattle sector



Project ID: 730924

Deliverable number: D5.1

Deliverable title: Optimised digestion trial protocols

EC version : V2

Due date of milestone	31/01/2022 (M48)
Actual submission date	17/06/2022 (M52)

DOCUMENT INFO

1. Author(s)

Organisation name lead contractor	INRAE
-----------------------------------	-------

Author	Organisation	e-mail
Chris Reynolds	URead	c.k.reynolds@reading.ac.uk
Zoe Barker	Uread	z.e.barker@reading.ac.uk
Dave Humphries	Uread	d.j.humphries@reading.ac.uk
Jan Dijkstra	WU	j.dijkstra@wu.ac.nl
Rene Baumont	INRAE	Rene.baumont@inrae.fr
Gonzalo Cantalapiedra	INRAE	gonzalo.cantalapiedra@inrae.fr
Peter Lund	AU	peter.lund@anis.au.dk
Björn Kuhla	FBN	b.kuhla@fhn-dummerstorf.de

2. Revision history

Version	Date	Modified by	Comments
V1	29 May 2022		
V2	13 September 2022	Chris Reynolds	To take into account EC Review remarks

3. Dissemination level

PU	Public	<input checked="" type="checkbox"/>
CO	Confidential , only for members of the consortium (including the Commission Services)	<input type="checkbox"/>

EXECUTIVE SUMMARY

Background	<p>In vivo techniques relying on total collection of faeces and urine (total tract diet digestion, N balance) are considered a Gold Standard Method (GSM) for phenotyping animals for feed efficiency (Martin, 1966; Spanghero and Kowalski, 1997), although the accuracy and precision of the results are compromised by technical limitations and there are concerns about staff, time, and safety, as well as animal welfare. Although the methods have been used for over 150 years, there is concern that aspects of their use at individual research facilities varies considerably, especially in terms of the number of days of collection used and the rigour in which collection of excreta and processing of samples is undertaken.</p> <p>One of the aims of SmartCow (https://www.smartcow.eu/) is to identify sources of variation in key GSM <i>in vivo</i> measurements of feed efficiency, including diet digestion and N excretion and balance. This will enable application of the 3-Rs by reducing variability, increasing precision and accuracy, and standardizing the methods and procedures used across research facilities within SmartCow installations and more widely. SmartCow WP5, working alongside WP3 for standardization issues, will evaluate and enhance the techniques being used, as well as contribute to the development of alternative approaches within WP6 through sample provision.</p> <p>As measurement of diet digestion is required for measurements of diet N utilization and balance, and measurements of faecal N excretion incorporate measurement of faecal dry matter excretion and digestion, it is difficult to separate measurements of ‘digestion’ and ‘N balance’ in the evaluation of historical data and techniques. For this reason, there is some unavoidable overlap in the deliverable reports 5.1 (optimized digestion trial protocols) and 5.2 (optimized N balance procedures).</p> <p>However, we differentiate our final recommendations between both documents, focussing on digestion protocols in D5.1 and more specifically on N balance procedures in D5.2.</p>
Objectives	<p>The overall aim is to identify and address sources of variation in key <i>in vivo</i> measurements of dietary nutrient use efficiency and associated emissions of nitrogen by cattle, thereby improving the measurements and unifying the approaches used across SmartCow installations, providing a global standard.</p> <p>The objectives are to (i) improve the accuracy and precision of measurements, and (ii) unify the methods used across SmartCow infrastructures.</p>

	Specific objectives are to develop optimised procedures for measurements of diet digestion using total collection of faeces that are acceptable in terms of precision, accuracy, animal welfare, and resource efficiency.
Methods	<p>We used 3 approaches:</p> <p>Meta-analysis: A meta-analysis of existing measurements of diet digestion and N balance for lactating dairy and growing beef cattle was conducted to determine the extent to which differences in measurement techniques across locations (e.g. collection methods, sample handling, days of sampling, etc.) introduces variation in measurements. Briefly, a database containing 2835 individual cow measurements of diet digestion and full N balance from digestion trials carried out at 12 research sites was assembled that include data from SmartCow partners Individual cow measurements included dry matter intake (DMI), chemical composition of the diets fed, nutrient intakes, faecal DM and N outputs, urinary N excretion, milk yield and composition, and body weight and other animal characteristics (e.g. breed, physiological state, sex). Multivariable analysis was used to account for variation due to experiment, diet intake, diet composition, and animal characteristics in order to ascertain the extent to which location and methodology accounted for variation in measurements of faecal excretion of DM and N.</p> <p>Equipment development: Equipment and techniques for collection and sampling of faeces and urine were improved and further developed as needed at individual SmartCow installations to reduce variability, improve precision, and improve animal health and welfare during measurements. This included design and development of new digestion stalls at Uread and INRAE. The results of equipment development are presented only in D5.1.</p> <p>Assessments of sources of variation: Sources of variation in digestion and N balance measurements used at Uread, INRAE, WU, FBN and AU were assessed according to the methods being used. Daily measurements of diet dry matter and N digestion (and urine N excretion) were obtained over the course of 4 to 10-day digestion trials conducted at Uread (lactating dairy cows) and INRAE (growing cattle) to determine the impact of number of days of excreta collection on measurements of diet digestion (and N excretion and balance). The data were also used to calculate the impact of days of collection on variation of the measurements and thus the number of animals required for measuring detectable differences. Sources of variation in digestion and N balance measurements were also evaluated by adding additional measurements to ongoing trials conducted at AU, FBN and WU. These results are detailed only in D5.2. Additional samples of faeces, urine, milk, and blood were obtained from some of these trials for WP6 proxy measurements.</p>

<p>Results & implications</p>	<p>The results of the digestion and N balance trial procedure evaluations (MS5.4) were reported and discussed in detail at a workshop held virtually on 5 January 2022. Minutes from the meeting and individual presentations are available on the SmartCow collaborative website (https://www.smartcow.eu/) and available on request. These presentations and the minutes of discussions highlighted improvements made at each SmartCow installation and identified key sources of variation and areas of focus for future improvement in the precision and accuracy of measurements of digestion and N balance in cattle.</p> <p>The meta-analysis of faecal DM and N and other N balance measurements found that for all data sets and variables, over half the variation in measured faecal DM and N excretion and N balance (retention) was explained by the effect of research site and individual experiment. Variation due to research site was less for faecal DM and N excretion (and digestion), but substantial variation due to experiment was observed (56 – 62 %), partly reflecting differences in experimental diets. There were no significant effects of specific methodology variables (e.g. days of collection or method of N analysis), thus the analysis suggests that variation across sites is due more to exactly how specific experimental methods are used, rather than the method used per se. In this regard, strict attention to detail in the conduct of methods and procedures and training of staff and students undertaking the experimental procedures at all levels, from feeding and collection of samples, through to sample storage, processing and analysis is critical for minimizing variation in measurements of digestion in cattle.</p> <p>At both INRAE and Uread, new stalls for housing animals during digestion measurements were developed that sought to improve animal comfort and welfare, health and safety of staff caring for the animals and obtaining samples, and provide flexibility for use with different sizes and types of animals. Key features included the ability to adjust the stalls for different sizes of animals and to temporarily restrict animal movements (e.g. adjustable side panels) during milking or sampling. A notable feature of the new stalls at Uread was the installation of waterbeds for stall matts, which provided a striking improvement in cow comfort.</p> <p>Key sources of variation in digestion trials addressed at individual locations included evaluation of effects of the number of days of sampling and the impact of spot sample collection timing. The SmartCow consortium Publisso publication ‘Methods in cattle physiology and behaviour research – Recommendations from the SmartCow consortium’ includes a chapter on ‘Nutrient Digestibility and Balance Studies’ that describes recommended procedures for conducting digestion trials and N balance (DOI: 10.5680/mcpb007). The results of the joint research activities for WP5 highlight key aspects of the procedures described in the book of methods and specific sources of variation that warrant further investigation, but do not suggest that major revisions of the current chapter are required. Some of the key considerations highlighted for reducing variance and increasing the precision and accuracy include:</p>
--	--

General:

1. Strict attention to detail and protocols is critical to minimizing variation in the measurements obtained. Improvements can be achieved through routine self-assessment and critical evaluation at each installation. These evaluations should be conducted at each experimental facility as the application of specific methods and procedures will vary due to the historical use of procedures and modifications that have occurred over time.

2. Training of staff and students conducting trials and effective communication is essential to be sure all are fully engaged and committed to the success of the experiments and are aware of the implications of specific procedures and sources of variation.

Specific procedures:

1. Under conditions of measurement at INRAE, repeatability of measurements of digestion in growing bulls were greater after at least 7 days of collection and for digestibility was greatest after 10 days of collection.

2. Shorter periods of 4 days were acceptable for lactating dairy cows in a trial where intake and diet composition did not vary over the course of the measurement periods.

3. Cow comfort and adaptation and training to digestion stalls and collection equipment and procedures is important for reducing variability and minimizing stress.

4. Accurate measurement of diet intake and composition of feeds and refusals is critical.

5. It is essential that collection of faeces and urine is complete and no cross contamination occurs.

6. If using markers and spot faecal samples for digestion trials sampling that characterizes more times of a day over fewer days is recommended over fewer times of day over more days. In the studies at AU 3 times per day over 2 days were recommended.

Table of content

1) Introduction	8
2) Objectives:.....	8
3) Methods:.....	8
4) Expected outcomes:.....	9
5) Key Results: Brief synopsis of the outcomes of the joint research activities for WP5:.....	10
6) Optimised digestion trial protocols (D5.1).....	15
7) References:	16
8) Appendix	18
SmartCow Summary of Results – N Balance Meta-analysis	18
Introduction	18
Methods	18
Database.....	18
Analysis of single explanatory variables	19
Multivariate Analysis	21
Results.....	21
Descriptive summaries	21
Retained N models	29
Urine N models.....	30
Faecal N models	30
Milk N models	32
Discussion	33
Conclusions	35
References	35

1) Introduction

Whilst *in vivo* measurements of diet digestion and N excretion and balance have been used for over 150 years and are considered to be 'standard' techniques, there is concern that the methods used at individual research facilities vary considerably, especially in terms of the number of days of collection used and the rigour in which collection of excreta and processing of samples is undertaken. Once all routes of N intake and excretion are measured, N balance is calculated as the difference and is assumed to represent total body tissue retention or loss of nitrogenous compounds. As reviewed previously, these measurements of body N accretion can often be excessively high relative to biologically realistic rates of body protein accretion and measured weight gain (Martin, 1966; Spanghero and Kowalski, 1997 and 2001; Reynolds and Kristensen, 2008; Hristov et al., 2019). As any potential losses of N from feed, faeces, urine, milk, claws, or as hair and scurf that are not measured are included in the estimate of body N balance, it has long been hypothesized that the sum of these volatile N (e.g. ammonia) and other unmeasured N losses may contribute to the excessively high rates of body N retention often reported (Spanghero and Kowalski, 1997). Recent studies have shown the extent to which sample processing and analytical method can affect measured N concentrations of feed, faeces, urine and milk (Morris et al., 2019). In addition, there has been a trend in the literature for the length of digestion trials to be reduced, which may in part be occurring to minimize weekend labour requirements and minimize cost of the experiment. Historically digestion trials have been conducted over 7 days or more to account for daily variation in faecal excretion (Schneider and Flatt, 1975), and it is not certain if there is a minimal number of days of collection required for a sufficient reduction in variance to be reached in order to minimize the number of animals required for statistically meaningful results. This may in part depend on the production level of the animal and the type of diet fed, as it has previously been shown the CV for faecal dry matter excretion is higher for lactating cows than for dry cows (Schneider and Flatt, 1975).

Measurement of diet digestion is required for measurements of diet N utilization and balance, and measurements of faecal N excretion require measurement of faecal dry matter excretion and digestion, therefore it is difficult to separate measurements of 'digestion' and 'N balance' in the evaluation of historical data and techniques. For this reason, there is some overlap between the deliverable reports 5.1 (optimized digestion trial protocols) and 5.2 (optimized N balance procedures) in the report that follows.

2) Objectives:

The overall objective of task 5.1 was to optimise procedures for measurements of 1) diet digestion and 2) nitrogen (N) balance using total collection of faeces and urine that are acceptable in terms of precision, accuracy, animal welfare, and resource efficiency and are applicable for lactating dairy and growing beef cattle.

3) Methods:

3.1) Meta-analysis of existing data: First, historical measurements of diet digestion (and N balance) for lactating dairy and growing beef cattle were compared through a meta-analysis to determine the extent to which differences in measurement techniques across locations (e.g. collection methods, sample handling, days of sampling, etc.) introduces variation and bias in measurements.

Methodology used and specific details and results are described in the associated report (MS5.1 – appendix 1). Briefly, a database containing 3969 individual cow measurements of diet digestion and

N excretion and balance from digestion trials carried out at 15 research sites associated with 11 different research institutions. The data base was assembled using data from SmartCow (<https://www.smartcow.eu/>) partners as well as the Feed and Nutrition Network of the Global Research Alliance (<https://globalresearchalliance.org/research/livestock/networks/feed-nutrition-network/>). Research sites were located in North America (3), Europe (11) and Oceania (1). The data set was comprised of 125 separate experiments with sites contributing data from 1 to 34 experiments (18 to 759 observations). Individual cow measurements included dry matter intake (DMI), chemical composition of the diets fed, nutrient intakes, faecal DM and N outputs, urinary N excretion, milk yield and composition, and body weight and other animal characteristics (e.g. breed, physiological state, sex). Retained N balance was calculated by subtracting faecal N, urine N and milk N (if lactating animals) excretions from N intake. Complete N balance data was available for 12 sites (2835 observations) and only observations associated with complete N balance (as opposed to only N digestion) measurement were included in the meta-analysis, which was conducted much as described previously (van Lingen et al., 2019) to account for variation due to experiment, diet, intake, diet composition, and animal characteristics in order to ascertain the extent to which location and methodology accounted for variation in measurements of faecal, urine and milk excretions of N and resulting N balance.

3.2) Equipment development: Secondly, equipment and techniques for collection and sampling of faeces and urine were improved and further developed as needed at individual SmartCow installations to reduce variability, improve precision, and improve animal health and welfare during measurements. This included design and development of new digestion stalls at Uread and INRAE.

3.3) Assessments of sources of variation: Thirdly, daily measurements of diet dry matter (and N digestion and urine N excretion) were obtained over the course of 4 to 10-day digestion trials conducted at Uread (lactating dairy cows) and INRAE (growing beef cattle) to determine the impact of number of days of excreta collection on measurements of diet digestion (and N excretion and balance). The data were also used to calculate the impact of days of collection on variation of the measurements and thus the number of animals required for detecting significant differences. Diets with low and excessive N concentration were used to produce a large difference in urine urea concentration and thus potential losses of ammonia from urine. Sources of variation in digestion and N balance measurements were also evaluated by adding additional measurements to ongoing trials conducted at AU, FBN and WU. Additional 'spot' samples of faeces, urine, milk, and blood were obtained from some of these trials for WP6 proxy measurements.

4) Expected outcomes:

Recommended measurement protocols that reduce variability of diet digestion measurement and minimise the number of animals and samples required from the animals used.

5) Key Results: Brief synopsis of the outcomes of the joint research activities for WP5:

The results of the digestion and N balance trial procedure evaluations (MS5.4) were reported and discussed in detail at a workshop held virtually on 5 January 2022. Minutes from the meeting and individual presentations are available on the SmartCow collaborative website or by request. These presentations and the minuted discussions highlighted improvements made at each SmartCow installation and identified key sources of variation and areas of focus for future improvement in the precision and accuracy of measurements of digestion and N balance in cattle.

5.1) Meta-analysis of Individual N Balance Measurements: A full report of the methodology used, results, and discussion of the meta-analysis of variation in measurements of faecal dry matter (linked to diet digestion) and faecal, urine, and milk N excretion and resulting N balance can be found in the full meta-analysis report (MS5.1; see below). Factors contributing to variation were assessed for measurements obtained in growing, nonlactating, and lactating cattle ($n = 2817$ or $n = 1704$) and lactating cattle only ($n = 1277$). Dietary and animal factors were addressed in order to better assess variation due to research site and methodology used. For all data sets and response variables over half the variation in measured N excretion and N balance (retention) was explained by the effect of the research site and individual experiment. Site accounted for from 20 to 27% of variation in urine N excretion and N balance, and 37% of milk N excretion (likely reflecting differences in milk yield of cows used at each site). Variation due to research site was less for faecal dry matter and N excretion (and digestion), but substantial variation due to experiment was observed (56 – 62%, partly reflecting differences in experimental diets). Much of the effect of experiment on faecal dry matter excretion in the all cattle evaluation appeared due to animal type, as the variation due to experiment was much lower when only data from lactating dairy cows was evaluated (41.2 vs 8.3%, respectively). Graphical representation of the variation in faecal dry matter and N excretion across research sites is shown in figures 1 and 2, respectively.




Figure 1. Relationship between faecal dry matter excretion and dry matter intake (kg/d) for individual research sites.




Figure 2. Relationship between N balance and N intake for individual research sites.

Variation in faecal dry matter and N excretion due to site was low relative to other component measurements of N balance (15.0 and 13.5 % for all cattle and lactating cattle models, respectively). There were no significant effects of specific methodology variables (e.g. days of collection or method of faecal excretion measurement), which suggests that apart from differences in the type of experimental animals studied, variation between sites was relatively low and may relate more to how the specific experimental methods are used, rather than the method used per se. In this regard strict attention to detail in the conduct of methods and procedures and training of staff and students undertaking the experimental procedures at all levels, from feeding and collection of samples, through to sample storage, processing and analysis is critical for minimizing variation in measurements of diet digestion (Snedecor and Flatt, 1970).

5.2) Equipment and procedure development:

At both INRAE and Uread new stalls for housing animals during digestion measurements were developed that sought to improve animal comfort and welfare, health and safety of staff caring for the animals and obtaining samples, and provide flexibility for use with different sizes and types of animals. Key features included the ability to adjust the stalls for different sizes of animals and to temporarily restrict animal movements (e.g. adjustable side panels) during milking or sampling. Pictures of the stalls at INRAE and Uread are presented in figures 3 and 4. A notable feature of the the new stalls at Uread was the installation of waterbeds for stall matts (Spinder Dairy Housing Concepts), which provided a striking improvement in cow comfort whilst lactating dairy cows were housed in the stalls. This led to noticeable improvements in intake and milk yield stability during measurements.



Figure 3. Digestion stall at INRAE.



Figure 4. Digestion stall at Uread.

5.3) Assessments of sources of variation:

As noted above, some of the key sources of variation in digestion trials addressed at individual locations included days of sampling and the impact of spot sample collection timing.

a) Use of markers and spot sampling (AU)

At **AU** markers and spot-sampling of faeces is used as an alternative to total faecal collection and a trial was conducted to determine the most appropriate schedule of spot-sample collection. It was concluded that 3 samplings (morning, early afternoon, late afternoon) over 2 consecutive days was most appropriate based on the variation observed (previously 2 samples over 3 days was used).

b) Effect of sample drying and grinding (FBN) (results presented in D5.2)

c) N losses during measurements of N balance (WU) (results presented in D5.2)

d) Assessment of the 'gold standard' measurement with 2 crude protein concentrations (UREAD)

Uread conducted a 4 x 3 switch-back design study using 4 mid-lactation Holstein cows fed ad libitum one of 2 total-mixed ration (TMR) diets differing in total CP concentration (14 vs 18%) and with 3 five-week periods. Cows were randomly assigned to one of the two diets (2 cows each) in the first period and then diets were switched at the beginning of each subsequent period (14-18-14 vs 18-14-18). Digestion and N balance was measured over 8-day periods using total collection of faeces and urine whilst cows were housed in respiration chambers. Samples of feed, feed refusals, faeces, urine and milk were obtained daily and kept chilled until analyzed for dry matter and N concentration as soon as possible on the day of collection. Representative daily samples were also taken from 24-hour collections and added to a frozen bulk sample obtained over 4 to 8 days. These bulk samples were then frozen and analysed as soon as possible after completion of the trial (within 10 days).

A key feature of the planning for this trial was the involvement of all staff involved in weekly meetings to co-develop new equipment and standard operating procedures for digestion trials at Uread. This training and attention to detail were considered essential for precise and accurate measurements.

Over the course of the 8 days of collection both feed intake and N concentration were relatively stable, with minimal variation in feed DM intake and excretion when comparing data averaged over days 1-4 versus 1-8 (Tables 3 and 4). This may in part reflect both cow comfort as noted above and also training and acclimatisation of cows to the digestion stalls and chambers prior to the trials. When comparing data from 4-day and 8-day collections, there were some small differences in the absolute values for feed DM intake and faecal DM excretion, however, the differences were not substantial (Tables 1 and 2). As expected, as sample size was increased from 4 to 8 days of collection the average standard errors associated with dry matter intake and excretion were slightly (approximately 8%) lower when 8 days of collection were included in the analysis. This suggests that when animals are well adapted to digestion stalls and equipment and diet intake and composition is stable then measurements periods of 4 days would be acceptable for measuring feed dry matter and protein digestibility in lactating dairy cows.

Table 1. Diet digestion and urine excretion based on an average of 8 daily samples

	Diet CP, %			P <1	
	14	18	SEM	Diet	Period
Dry matter intake, kg/d	19.53	20.98	0.958	0.134	0.614
Milk yield, kg/d	26.4	28.8	1.24	0.292	0.048
Faecal dry matter, kg/d	6.08	6.30	0.306	0.571	0.669
Urine, kg/d	18.47	22.94	0.747	0.001	0.019
Dry matter digested, kg/d	13.32	14.64	0.713	0.116	0.641
Dry matter digested, g/g	0.686	0.698	0.009	0.118	0.711

Table 2. Diet digestion and urine excretion based on 4 day bulk sample analysis.

	Diet CP, %			P <	
	14	18	SEM	Diet	Period
Dry matter intake, kg/d	19.62	21.17	1.044	0.159	0.706
Faecal dry matter, kg/d	6.08	6.18	0.333	0.799	0.472
Urine, kg/d	18.46	22.75	0.921	0.003	0.033
Dry matter digested, kg/d	13.42	14.94	0.761	0.105	0.791
Dry matter digested, g/g	0.695	0.702	0.009	0.377	0.776

e) Evaluation of the effect of days of collection on the repeatability and precision of measurements (INRAE)

INRAE conducted an extensive evaluation of the effect of days of collection on the repeatability and precision of measurements of feed digestion (and N excretion and balance) of 16 growing bulls fed ad libitum diets differing in CP concentrations (17.3 vs 11.6 % CP, 8 bulls per diet) over the course of 2 digestion trials repeated at 9 week intervals. Bulls remained on the same dietary treatment through both digestion trials. The bulls were moved from free-stall housing to digestion stalls for 15 days, with 5 days of adaptation followed by 10 days of sample collection. Hair cortisol was also measured as an indicator of animal stress. The results showed that repeatability, measured as the within-diet correlation between measurements in the 2 periods, increased with the length of the collection periods for feed dry matter digestibility. In addition, power tests were conducted to determine the effect of days of collection on the minimum detectable difference for digestibility. The results show that longer periods result in a lower detectable difference for dry matter digestibility (Figure 6), with the lowest detectable difference for 10 day collections.. Measurements of hair cortisol were lowest when animals were housed in free stalls and increased during digestion trials, although concentrations were highest in the first days of collection, suggesting that animals adapted to the stalls over time.




Figure 6. Minimum detectable difference (MDD, g/100g) and days of measurement of feed dry matter digestibility for 1, 4, 8 and 16 growing bulls.

6) Optimised digestion trial protocols (D5.1).

Collective recommendations for best practice procedures for the conduct of nutrient digestibility were published by the SmartCow consortium in the Publisso publication 'Methods in cattle physiology and behaviour research – Recommendations from the SmartCow consortium'. The chapter 'Nutrient Digestibility and Balance Studies' describes recommended procedures for conducting digestion trials and N balance.

Danesh Mesgaran S, Kuhla B, Baumont R, Cantalapiedra-Hijar G, Nozière P, Lund P, Humphries D, Dijkstra J. Nutrient digestibility and balance studies. In: Mesgaran SD, Baumont R, Munksgaard L, Humphries D, Kennedy E, Dijkstra J, Dewhurst R, Ferguson H, Terré M, Kuhla B, (editors). Methods in cattle physiology and behaviour – Recommendations from the SmartCow consortium. Cologne: PUBLISSO; 2020. DOI: 10.5680/mcpb007 (https://books.publisso.de/en/publisso_gold/publishing/books/overview/53/188).

The results of the joint research activities for WP5 highlight key aspects of the procedures described, but do not suggest that major revisions of the current chapter are required. Some of the key considerations for reducing variance and increasing the minimum detectable difference include:

General:

1. Strict attention to detail and protocols is critical to minimizing variation in the measurements obtained. Improvements can be achieved through routine self assessment and critical evaluation at each installation. These evaluations should be conducted at each experimental facility as the application of specific methods and procedures will vary due to the historical use of procedures and modifications that have occurred over time.

2. Training of staff and students conducting trials and effective communication is essential to be sure all are fully engaged and committed to the success of the experiments and are aware of the implications of specific procedures and sources of variation.

Specific procedures:

1. Under conditions of measurement at INRAE repeatability of measurements of digestion (and N balance) of growing beef cattle were greater after at least 7 days of collection and for digestibility was greatest after 10 days of collection.
2. Shorter periods of 4 days were acceptable in a trial with lactating dairy cows at Reading where cows were trained to digestion stalls before experiments began and intake and diet composition did not vary over the course of measurement periods. The current EU directive is for collection periods to not exceed 5 days although a derogation can be obtained if justified.
3. Cow comfort and adaptation and training to digestion stalls and collection equipment and procedures is important for reducing variability and minimizing stress.
4. Accurate measurement of diet intake and composition of feeds and refusals is critical.
- 5.

If using markers and spot faecal samples for digestion trials sampling that characterizes more times of a day over fewer days is recommended over fewer times of day over more days. In the studies at AU 3 times per day over 2 days were recommended.

7) References:

Hristov, A. N., A. Bannink, L. A. Crompton, P. Huhtanen, M. Kreuzer, M. McGee, P. Nozière, C. K. Reynolds, A. R. Bayat, D. R. Yáñez-Ruiz, J. Dijkstra, E. Kebreab, A. Schwarm, K. J. Shingfield, Z. Yu. Nitrogen in ruminant nutrition: a review of measurement techniques. *Journal of Dairy Science*, 2019, 102, 5811-5852. 10.3168/jds.2018-15829

Martin, A. K. 1966. Some errors in the determination of nitrogen retention of sheep by nitrogen balance studies. *Br. J. Nutr.* 20:325-337.

Morris, D. L., A. W. Tebbe, W. P. Weiss, and C. Lee. 2019. Short communication: Effects of drying and analytical methods on nitrogen concentrations of feeds, feces, milk, and urine of dairy cows. *J. Dairy Sci.* 102:5212-5218.

Reynolds, C. K., and N. B. Kristensen. 2008. Nitrogen recycling through the gut and the nitrogen economy of ruminants: an asynchronous symbiosis. *J. Anim. Sci.* 86:E293-E305.

Schneider, B. H., and W. P. Flatt. 1975. *The Evaluation of Feeds through Digestibility Experiments.* The University of Georgia Press, Athens, GA.

Spanghero, M., and Z. M. Kowalski. 1997. Critical analysis of N balance experiments with lactating cows. *Livest. Prod. Sci.* 52:113-122.

Spanghero, M., and Z. M. Kowalski. 2021. Updating analysis of nitrogen balance experiments in dairy cows. *J. of Dairy Sci.* Volume 104:7725-7737. <https://doi.org/10.3168/jds.2020-19656>.

van Lingen, H.J., Niu, M., Kebreab, E., Valadares Filho, S.C., Rooke, J.A., Duthie, C.-A., Schwarm, A., Kreuzer, M., Hynd, P.I., Caetano, M., Eugène, M., Martin, C., McGee, M., O'Kiely, P., Hünerberg, M., McAllister, T.A., Berchielli, T.T., Messana, J.D., Peiren, N., Chaves, A.V., Charmley, E., Cole, N.A.,

Hales, K.E., Lee, S.-S., Berndt, A., Reynolds, C.K., Crompton, L.A., Bayat, A.-R., Yáñez-Ruiz, D.R., Yu, Z., Bannink, A., Dijkstra, J., Casper, D.P., Hristov, A.N., 2019. Prediction of enteric methane production, yield and intensity of beef cattle using an intercontinental database. *Agric. Ecosyst. Environ.* 283, 106575. <https://doi.org/10.1016/j.agee.2019.106575>

8) Appendix

8.1) Appendix 1: MS5.1 Evaluation of historical data for variation in digestion and N balance data.

SmartCow Summary of Results – N Balance Meta-analysis

Introduction

Measurements of feed efficiency related to a ‘sustainable phenotype’ for cattle include feed digestion, nitrogen (N) excretion in faeces, urine, and milk, and N balance. However, it is important to ‘refine’ the techniques used for these measurements in order to minimize technique associated variation and thereby ‘reduce’ the number of animals required for our experiments when alternative approaches such as proxies and biomarkers (WP6) are not appropriate (Hristov et al., 2019). Moreover, from an environmental perspective, accurate estimation of N retention is of large importance because dietary N consumed and not retained in animal tissues or secreted in milk is excreted in urine and faeces. This contributes to water pollution, gaseous N emissions including ammonia and nitrous oxide, and small particulate matter formation in the atmosphere (Bougouin et al., 2022). Urinary N is much more labile and susceptible to fast leaching and volatilization losses than faecal N, and therefore accurate estimation of faecal vs urine N excretion is required. Thus key measurements are whole digestive tract diet digestion and associated measurements of N excretion in faeces and urine required (with milk N excretion measurement if needed) to measure whole body N utilization, typically based on total collection of urine and faeces, with separation of faeces and urine maintained to minimize volatile N losses during collection (Schneider and Flatt, 1975; Hristov et al., 2019). Whilst these measurements have a long history of use, there is wide variation in the standard operating procedures in use at individual locations and it has long been known that measurements of N balance are often subject to large errors of measurement (Hristov et al., 2019; Martin, 1966; Sphanghero and Kowalski, 1997 and 2021). Known sources of measurement error include cross-contamination of faeces and urine, which can lead to urea catabolism and ammonia loss, N losses during sample preparation, storage, drying, and grinding prior to analysis, insufficient mixing of urine and acid added during collection to minimize ammonia loss, and the number of days of collection. Historically collection periods of 10 days or more were used for digestion trials in dairy cattle to account for daily variations in faecal output, whilst more recently collection period of 4 days have become common, as this negates the need for weekend work.

Therefore, our objective was to assemble a data base of N balance measurements for cattle and associated diet and production variables and conduct a meta-analysis to determine the extent to which variation in measurements is attributable to research location and the methods used, after accounting for variation due to feed DMI and composition and other animal characteristics that may affect N excretion and tissue balance. This will provide an evidence base for recommendations of ‘best practice’ at SmartCow research facilities that increase precision and accuracy and minimize animal numbers required for ‘in vivo’ measurements of N balance.

Methods

Database

The complete data set was comprised of 3969 individual cow measurements of diet digestion and N excretion and balance from digestion trials carried out at 15 research sites associated with 11 different research institutions. The data base was assembled using data from SmartCow

(<https://www.smartcow.eu/>) partners as well as participants in the Food and Nutrition Network of the Global Alliance (<https://globalresearchalliance.org/research/livestock/networks/feed-nutrition-network/>). Research sites were located in North America (3), Europe (11) and Oceania (1). The data set was comprised of 125 separate experiments with sites contributing data from 1 to 34 experiments (18 to 759 observations). Individual cow measurements included dry matter intake (DMI), chemical composition of the diets fed, nutrient intakes, faecal DM and N outputs, urinary N excretion, milk yield and composition, and body weight and other animal characteristics (e.g. breed, physiological state, sex). Retained N balance was calculated by subtracting faecal N, urine N and milk N (if lactating animals) excretions from N intake. Complete N balance data was available for 12 sites allowing the calculation of N balance for these observations (2835 observations). Figure 1 summarises the entire data set and subsequent subsets of data used in the data analysis processes.




Figure 1 – Summary data sets considered in the modelling processes with details of data exclusions made.

Analysis of single explanatory variables

Variables with a high proportion of missing observations (>60% of the total number of individual cow measurements) were excluded from models used in further analyses. It was considered important that dietary N intake was accounted for in the models as the key determinant of N excretion in faeces, urine and milk. To achieve this, each response variable (retained N, urine N excretion, faecal N excretion, milk N) was tested with either: 1) N intake or 2) DMI and CP content as explanatory variables. The model fit for the N intake models was better based on AIC and BIC values, so NI was

included in all subsequent models of single explanatory variables. First, a correlation matrix was produced for numerical variables to identify highly correlated variables, $|r| = 0.75$ (Appendix 1). DMI and N intake were highly correlated to a number of other dietary intake measures, as DMI determined the total intake of all diet components. Therefore, the decision was made to take forward N intake and diet ash, NDF, ADF and starch concentrations for further analysis along with a range of methodology-related variables. Methodological variable categories were: treatment type (diet change, feed additive, abomasal infusion, delivery method, cow status), experimental design (change over, randomised), DMI measurement method (weighed manually or electronically), feed presentation (total mixed ration, separate forage and concentrate, pellets, cut and carry), and collection method (total collection, manure – faecal N, faecal marker and spot collection of urine, faecal marker with no urine collection). This reduced selection of relevant diet composition, animal characteristic, and methodological variables (Table 1) were tested individually in a series of mixed models where retained N, urine N excretion, faecal N excretion or milk N excretion were the outcome variables. Experiment nested within research site were random effects unless there was insufficient variation to support nesting experiment within site. Variables were selected for further model development if their effect was significant or had a nonsignificant trend ($P < 0.10$). Table 1 summarises the results of the single variable (plus N intake) analysis and shows the short list of variables considered for further analysis. Methodological variables (Table 2) included treatment type (e.g. diet change or feed additive), experimental design (continuous or change over), duration of adaptation to treatment, days of collection, N analysis method (Kjeldahl or combustion), DMI measurement method (manual or automated weighing), intake level (ad lib vs restricted), feed presentation (e.g. total mixed ration [TMR], pellets, etc.), and collection method (e.g. total or marker).

Table 1. Summary of single variable analysis where ✓ = $p < 0.05$, $p = 0.1-0.05$ = NST and × = $p \geq 0.1$

	Retained N	Urine N excretion	Faecal N excretion ¹	Milk N excretion (Dairy only subset)
N intake	✓	✓	✓	✓
Cow status	✓	✓	✓	NA
Treatment type	×	✓	×	×
Experimental design	×	✓	✓	×
Adaptation duration	×	×	×	×
Duration of collection	×	×	×	NST
N analysis method	NST	×	NST	✓ ¹
DMI Measurement method	NST	✓	×	✓
Feed restriction	✓	✓	×	×
Feed presentation	×	✓	×	✓
Collection method	×	×	×	×
Lactation	✓	✓	✓	✓
Sex	×	×	×	NA
Body weight	✓	✓	×	✓
Ash content	✓	✓	✓	✓
NDF content	×	×	✓	✓
ADF content	×	×	✓	✓
Diet forage proportion	×	×	×	✓

¹ random effect of experiment used as insufficient variation to support the nesting of experiment within site

Multivariate Analysis

Outcome variables were retained N (N balance), urine N excretion, faecal N excretion or milk N excretion. Each were tested in a linear mixed models with experiment nested within site as random effects with random intercepts. An automated model build function based on lowest AIC while controlling for variance inflation factors was used to determine the 'best model' fit using R version 4.0.3 (R Core Team, 2020; van Lingen et al., 2019). Variables from table 1 that were $p < 0.10$ for one of more of the outcome variables were taken forward for the model build process. Highly correlated variables identified in the correlation matrix i.e. ADF content and NDF content were prevented from appearing in the model at the same time as each other. The inclusion of experiment duration and N chemistry method (i.e. Dumas or Kjeldahl) in combination with other variables resulted in high VIF. As experiment duration was non-significant for retained N, urine N and faecal N and only resulted in a non-significant trend for milk N it was decided to include N chemistry method in the short list of variables for the model build in which all combinations of the following variables were tested: N intake, cow status, body weight, diet ASH content, diet NDF content, diet ADF content, experiment design, DMI method, N chemistry method, feed offered (e.g. ad lib or restricted), collection method, treatment type, feed presentation, and forage proportion. Parity was only included in the short list for the milk N models. The sum of the variance associated with each random effect (site and experiment) and the residual variance were used to calculate a percentage variance associated with each.

Results

Descriptive summaries

The complete data set includes lactating beef cattle, growing cattle and dry dairy cattle but experiments with lactating dairy cattle dominate the data set and as a result the majority (85%) of animals are multiparous females (table 2). After lactating dairy, growing cattle was the next largest animal status category accounting for 10% of the total data set. However, many of the animal and methodology variables contained categories with no or very few data points reported, precluding the use of the growing animal only data set to determine causes of variation in the measurement of N balance. Beef breeds were not requested in the original data gathering exercise, so it is not possible to determine differences between other non-specified dairy breeds and beef breeds. The methodologies used in the experiments are summarised in table 2. Latin square and randomised experimental designs were dominant as was total collection of urine and faeces.

Table 2. Summary of categorical data

Categories	Entire data set n (%)	Complete N balance n (%)	Complete N Balance – Lactating Dairy n (%)	Complete N Balance – Growing n (%)
Animals				
Cow Status				
Dry dairy	160 (4.0)	160 (5.6)	0 (0.0)	0 (0.0)
Growing	408 (10.3)	308 (10.9)	0 (0.0)	308 (100.0)
Lactating beef	34 (0.9)	34 (1.2)	0 (0.0)	0 (0.0)
Lactating dairy	3367 (84.8)	2333 (82.3)	2333 (100.0)	0 (0.0)
Breed				
Holstein Friesian	2897 (73.0)	2019 (71.2)	1795 (76.9)	91 (29.5)

Jersey	58 (1.5)	58 (2.0)	42 (1.8)	0 (0.0)
Ayrshire	112 (2.8)	88 (3.1)	88 (3.8)	0 (0.0)
Brown Swiss	245 (6.2)	132 (4.7)	120 (5.1)	12 (3.9)
HF x Jersey	12 (0.3)	12 (0.4)	12 (0.5)	0 (0.0)
Other	373 (9.4)	277 (9.8)	27 (1.2)	205 (66.6)
Unknown	272 (6.9)	249 (8.8)	249 (10.7)	0 (0.0)
<i>Sex</i>				
Male	187 (4.7)	91 (3.2)	0 (0.0)	91 (29.5)
Female	3587 (90.4)	2549 (89.9)	2333 (100.0)	56 (18.2)
Unknown	195 (4.9)	195 (6.9)	0 (0.0)	161 (52.3)
<i>Lactation category</i>				
Multiparous	2185 (55.1)	1582 (55.8)	1482 (63.5)	0 (0.0)
Not-applicable	468 (11.8)	368 (13.0)	0 (0.0)	308 (100.0)
Primiparous	552 (13.9)	302 (10.7)	290 (12.4)	0 (0.0)
Unknown	764 (19.2)	583 (20.6)	561 (24.0)	0 (0.0)
Methodologies				
<i>Type of experiment</i>				
Change over	2745 (69.28)	1809 (63.8)	1481 (63.5)	280 (90.9)
Randomised trial	1204 (30.3)	1006 (35.5)	832 (35.7)	28 (9.1)
Unknown	20 (0.5)	20 (0.7)	20 (0.9)	0 (0.0)
<i>Duration of adaptation</i>				
0-6 days	61 (1.5)	60 (2.1)	60 (2.6)	0 (0.0)
7-15 days	1117 (28.1)	832 (29.3)	801 (34.3)	12 (3.9)
Over 15 days	1352 (34.1)	596 (21.0)	564 (24.2)	0 (0.0)
unknown	1436 (36.3)	1347 (47.5)	908 (38.9)	296 (96.1)
<i>Duration of collection</i>				
2 days	108 (2.7)	72 (2.5)	72 (3.1)	0 (0.0)
3 days	93 (2.3)	92 (3.2)	92 (3.9)	0 (0.0)
4 days	782 (19.7)	137 (4.8)	137 (5.9)	0 (0.0)
5 days	928 (23.4)	856 (30.2)	729 (31.2)	44 (14.3)
6 days	564 (14.2)	506 (17.8)	495 (21.2)	0 (0.0)
7 days	976 (24.6)	866 (30.5)	522 (22.4)	252 (81.8)
8 days	370 (9.3)	158 (5.6)	138 (5.9)	12 (3.9)
Unknown	148 (3.7)	148 (5.2)	148 (6.3)	0 (0.0)
<i>Intake method</i>				
Weighed electronically	900 (22.7)	140 (4.9)	140 (6.0)	0 (0.0)
Weighed manually	3069 (77.3)	2695 (95.1)	2193 (94.0)	308 (100.0)
<i>Wet chemistry methods</i>				
Dumas All	1620 (40.8)	746 (26.3)	710 (30.4)	12 (3.9)
Kjeldahl All	1867 (47.0)	1663 (58.7)	1229 (52.7)	296 (96.1)
Mixture	270 (6.8)	266 (9.4)	234 (10.0)	0 (0.0)
NIR	12 (0.3)	12 (0.4)	12 (0.5)	0 (0.0)
unknown	200 (5.0)	148 (5.2)	148 (6.3)	0 (0.0)
<i>Collection method</i>				
Faecal marker no urine	602 (15.2)	0 (0.0)	0 (0.0)	0 (0.0)
Faecal marker urine spot	162 (4.1)	72 (2.5)	72 (3.1)	0 (0.0)
Manure minus Faecal marker	214 (5.4)	158 (5.6)	158 (6.8)	0 (0.0)
Total collection	2991 (75.4)	2605 (91.9)	2103 (90.1)	308 (100.0)

A summary of diet composition, nutrient intake, diet digestibility and nitrogen outputs and retention variables are presented in table 3. Summaries of the additional data subsets were also produced but are not presented here as there were few discernible differences in mean values. The data were reviewed and checked for data points outside of expected ranges. Where potential errors were identified or where values were not considered biologically plausible, contributors were contacted to check their data sets and confirm any known reasons why the data should be excluded. As this aim of this meta-analysis was to consider sources of variation in the measurement of nitrogen excretion in cattle all data were retained unless there was a known error.


Table 3. Summary of diet composition, nutrient intake, diet digestibility and nitrogen outputs and retention for the entire study data set (3969 observations).

	Mean	SD	Min	Max	CV	n
Diet Composition						
CP concentration (g/kg DM)	163	26	71	435	0.16	3932
EE concentration (g/kg DM)	35	13	7	88	0.37	1919
ASH concentration (g/kg DM)	75	17	28	154	0.22	3367
NDF concentration (g/kg DM)	350	69	136	655	0.20	3455
ADF concentration (g/kg DM)	209	51	66	397	0.24	2575
Starch concentration (g/kg DM)	232	116	1	641	0.50	2471
Nutrient intakes						
DM intake (kg/d)	18.4	6.2	2.1	37.5	0.33	3962
Forage proportion (% diet DM)	60.1	16.4	4.7	100.0	0.27	3360
CP intake (kg/d)	3.1	1.1	0.3	7.2	0.37	3377
N intake (g/d)	493	180	43	11528	0.36	3735
EE intake (kg/d)	0.62	0.39	0.02	2.19	0.62	1692
ASH intake (kg/d)	1.4	0.5	0.1	4.6	0.38	3013
NDF intake (kg/d)	6.6	2.1	0.4	13.9	0.32	3165
ADF intake (kg/d)	3.8	1.5	0.2	9.5	0.39	2532
Starch intake (kg/d)	4.1	1.8	0.0	12.3	0.45	2257
Diet digestibility						
DM digestibility (%)	69.7	6.8	22.9	87.0	0.10	3142
OM digestibility (%)	71.8	4.5	52.4	88.4	0.06	2628
N digestibility (%)	66.7	6.8	27.4	88.5	0.10	3026
NDF digestibility (%)	55.5	10.9	4.4	91.2	0.20	2908
ADF digestibility (%)	52.2	12.3	5.0	92.6	0.24	2037
Starch digestibility (%)	95.8	4.6	43.4	100.0	0.05	1791
Nitrogen output and retention (using observations with complete N balance data sets only)						
Total N excretion (g/d)*	500	130	167	936	0.26	2333
Faecal N excretion (g/d)	156	65	11	385	0.41	2835
Urine N excretion (g/d)	159	73	4	446	0.46	2835
Milk N (g/d)*	151	42	41	296	0.28	2333
Retained N (g/d)	28	53	-279	315	1.94	2835
Faecal N as % of N intake	34.8	7.6	10.6	93.5	0.22	2835
Urine N as % of N intake	36.0	13.6	2.2	99.3	0.38	2835
Manure N as % of N intake	69.7	13.9	28.0	167.8	0.20	2835


Milk N as % of N intake*	29.2	6.8	9.3	65.7	0.23	2333
Retained N as % of N intake	6.1	13.4	-130.6	57.9	2.18	2835
Animal Data						
Body weight (kg)	612	126.3	129.0	974.0	0.21	3279
Milk production (kg/d)*	29.9	9.1	5.9	63.5	0.30	2310
Days in Milk (at exp start)*	137	78.7	4	567	0.61	1448

*Lactating dairy only


Variation associated with site and key methodologies are visualised in figures 2a-e (N balance), 3a-e (urine N), 4a-e (Faecal N), and 5a-e (Faecal dry matter).






Figures 2a-e. Data visualisations of N intake v N Balance plotted with regression lines for location and methodology classifications.






Figures 3a-e. Data visualisations of N intake vs. Urine N plotted with regression lines for location and methodology classifications.



Figures 4a-e. Data visualisations of N intake vs. Faecal N plotted with regression lines for location and methodology classifications.





Figures 5a-f. Data visualisations of dry matter intake vs. faecal dry matter plotted with regression lines for location and methodology classifications.

Retained N models

The best models for retained N for the maximal all animal data set (B), the all animal data set with exclusions (C) and the lactating cow only data set with exclusions (G) are presented in tables 4a-c respectively (see Figure 1). While the models presented in tables 4a and 4b contain the same variables the significance and effect sizes vary for the status variable. Where the model is produced for the lactating dairy cow only data set body weight and diet ash content are retained in the best model. Site level variation ranges from 21-27% across all models.

Table 4a. Best linear mixed effects model for retained N using data set B, maximum observations.

		Coefficient	SE	df	T-value	P value
<i>Intercept</i>		-18	10	15	-1.770	0.097
N intake		0.24	0.01	2813	26.925	<0.001
Status	Ref: dry dairy					
	Growing	25.24	15.85	119	1.592	0.114
	Lactating beef	-2.41	18.87	228	-0.127	0.899
	Lactating dairy	-82.04	4.88	2808	-16.814	<0.001

No. observations = 2817, No. of experiments = 106, No. of Sites = 11,
random effect = site|experiment,
Model variance: experiment = 37.9%, site = 21.1%, residual 40.9%

Table 4b. Best linear mixed effects model for retained N using data set C, with exclusions.

		Coefficient	SE	df	T-value	P value
<i>Intercept</i>		-29	12	19	-2.47	0.024
N intake		0.21	0.01	1698	19.32	<0.001
Status	Ref: dry dairy					
	Growing	49.88	13.91	86	3.59	<0.001
	Lactating beef	24.80	16.88	172	1.47	0.144
	Lactating dairy	-46.95	7.45	1475	-6.31	<0.001

No. observations = 1704, No. of experiments = 61, No. of Sites = 9,
random effect = site|experiment,
Model variance: experiment = 27.8%, site = 26.6%, residual 45.6%

Table 4c. Best linear mixed effects model for retained N using data set G, lactating dairy cows only with exclusions.

		Coefficient	SE	df	T-value	P value
<i>Intercept</i>		88	18	98	-4.856	<0.001
N intake		0.226	0.014	1263	16.370	<0.001
Body weight		-0.059	0.018	1275	-3.345	<0.001
Ash content		0.609	0.141	1167	4.323	<0.001

No. observations = 1277, No. of experiments = 49, No. of Sites = 9,

random effect = site|experiment

Model variance: experiment = 28.2%, site = 22.9%, residual 48.8%

Urine N models

The best models for urine N for the maximal all animal data set (B), the all animal data set with exclusions (C), and the lactating cow only data set with exclusions (G) are presented in tables 5a-c respectively (Figure 1). All models contain the same variables, and the results are highly consistent between models. Site level variation ranges from 20-25% across all models.

Table 5a. Best linear mixed effects model for urine N excretion using data set B, maximum observations.

	Coefficient	SE	df	T-value	P value
<i>Intercept</i>	-22	10	22	-2.10	0.048
N intake	0.26	0.01	2256	34.94	<0.001
Body weight	0.10	0.01	2272	9.73	<0.001

No. observations = 2274, No. of experiments = 82, No. of Sites = 11,

random effect = site|experiment,

Model variance: experiment = 42.7%, site = 20.3%, residual 37.0%

Table 5b. Best linear mixed effects model for urine N excretion using data set C, with exclusions

	Coefficient	SE	df	T-value	P value
<i>Intercept</i>	-26	11	20	-2	0.0313
N intake	0.26	0.01	1638	29	<0.001
Body weight	0.11	0.01	1703	10	<0.001

No. observations = 1704, No. of experiments = 61, No. of Sites = 9,

random effect = site|experiment,

Model variance: experiment = 35.5%, site = 23.1%, residual 41.4%

Table 5c. Best linear mixed effects model for urine N excretion using data set G, lactating dairy cows only with exclusions.

	Coefficient	SE	df	T-value	P value
<i>Intercept</i>	-35	14	28	-2.50	0.019
N intake	0.28	0.01	1276	22.84	<0.001
Body weight	0.10	0.02	1270	6.26	<0.001

No. observations = 1277, No. of experiments = 49, No. of Sites = 9,

random effect = site|experiment,

Model variance: experiment = 33.1%, site = 24.8%, residual 42.1%

Faecal N models

A lack of variation prevents the use of experiment nested within site with faecal N as the response variable. Therefore, the best models for faecal N for the maximal all animal data set (B), the all-animal data set with exclusions (C) and the lactating cow only data set with exclusions (G) are presented for a models with experiment as a random effect in tables 6a-c.

Table 6a. Best linear mixed effects model for faecal N excretion using data set B, maximum observations

		Coefficient	SE	df	T-value	P value
<i>Intercept</i>		16	4	351	4.39	<0.001
N intake		0.28	0.00	2817	56.69	<0.001
Status	Ref: dry dairy					
	Growing	-7.17	9.23	119	-0.78	0.439
	Lactating beef	-4.74	10.70	209	-0.44	0.658
	Lactating dairy	15.66	2.65	2813	5.90	<0.001

No. observations = 2817, No. of experiments = 106, random effect = experiment

Model variance: experiment = 62.4%, residual 37.6%

Table 6b. Best linear mixed effects model for Faecal N excretion – data set C, with exclusions

		Coefficient	SE	df	T-value	P value
<i>Intercept</i>		0	5	253	-0.03	0.974
N intake		0.27	0.01	1697	42.96	<0.001
Status	Ref: dry dairy					
	Growing	8.93	9.32	79	0.96	0.341
	Lactating beef	11.35	10.78	138	1.05	0.294
	Lactating dairy	30.17	4.39	1648	6.88	<0.001

No. observations = 1704, No. of experiments = 61, random effect = experiment

Model variance: experiment = 58.7%, residual 41.3%

Table 6c. Best linear mixed effects model for Faecal N excretion – data set G, lactating dairy cows only with exclusions

		Coefficient	SE	df	T-value	P value
<i>Intercept</i>		32	10	744	3.20	<0.001
N intake		0.28	0.01	1265	36.88	<0.001
Ash content		-0.38	0.08	1252	-4.51	<0.001
ADF content		0.11	0.03	1198	3.40	<0.001

No. observations = 1277, No. of experiments = 49, random effect = experiment

Model variance: experiment = 56.0%, residual 44.0%

Milk N models

The lactating cow only data set with exclusions (G) was used to produce a best model for milk N and is presented in table 7d. Parity was also retained in the best model for milk N with primiparous cows having lower milk N than multiparous cows. Site accounted for 34% of the variation.

Table 7d. Best linear mixed effects model for Milk N – data set G, lactating dairy cows only with exclusions

Explanations

		Coefficient	SE	df	T-value	P value
Intercept		121	12	74	10.44	<0.001
N intake		0.22	0.01	1271	30.67	<0.001
Body weight		-0.05	0.01	1268	-5.22	<0.001
Ash content		-0.50	0.07	1258	-6.79	<0.001
ADF content		-0.11	0.03	1149	-4.12	<0.001
Parity	Ref: Multiparous					
	Primiparous	-12.17	2.50	1164	-4.87	<0.001
	Unknown	9.77	5.53	533	1.77	0.078

No. observations = 1277, No. of experiments = 49, No. of Sites = 9,

random effect = site|experiment,

Model variance: experiment = 26.3%, site = 34.3%, residual 39.4%

Faecal dry matter excretion models

The best models for faecal DM for the all animal data set with exclusions (C) and the lactating cow only data set with exclusions (G) are presented in tables 8a-b respectively (see Figure 1). While the models presented in tables 8a and 8b contain the same variables the experiment variation accounts for a larger proportion of total variation with the all cattle model. This may relate to confounding between experiment and animal status (growing vs lactating cattle). For both models the variation due to site is relatively low.

Table 8a. Best linear mixed effects model for faecal DM using data set C, with exclusions.

	Coefficient	SE	df	T-value	P value
<i>Intercept</i>	-0.149	0.240	58	-0.624	0.535
Dry Matter Intake	0.334	0.004	1613	80.017	<0.001
Body weight	-0.001	0.000	1683	-5.175	<0.001
Forage %	-0.008	0.002	1584	-3.719	<0.001
CP concentration	-0.004	0.001	1618	-4.556	<0.001
ADF concentration	0.006	0.001	1622	10.369	<0.001

No. observations = 1686, No. of experiments = 57, No. of Sites = 7,

random effect = site|experiment,

Model variance: experiment = 41.6.0%, site = 15.0%, residual 43.4%

Table 8b. Best linear mixed effects model for faecal DM using data set G, lactating dairy cows only with exclusions.

	Coefficient	SE	df	T-value	P value
<i>Intercept</i>	0.438	0.392	229	1.117	0.265
Dry Matter Intake	0.322	0.006	1256	52.625	<0.001
Body weight	-0.001	0.000	1255	-4.164	<0.001
Forage %	-0.013	0.003	1119	-4.586	<0.001
CP concentration	-0.005	0.001	1180	-3.708	<0.001
ADF concentration	0.007	0.001	1173	8.561	<0.001

No. observations = 1279, No. of experiments = 47, No. of Sites = 7,

random effect = site|experiment

Model variance: experiment = 8.3%, site = 13.5%, residual 78.2%

Discussion

As observed in previous analyses (e.g. Angelidis et al., 2019; Bougouin et al., 2022), N intake was the major determinant of N excretion in faeces, urine and milk of cattle in the present study. When data from growing and nonlactating cattle are included in the data analysed, cattle type also had an effect on N excretion in faeces and N retention, with greater relative amounts of faecal N excretion and less body N retention for lactating dairy cows compared to growing, dry and lactating beef cattle. For faecal N excretion this may reflect effects of higher levels of DMI and diets with higher digestibility that result in greater faecal outputs, including greater endogenous faecal N associated with increased hindgut fermentation of carbohydrates in lactating dairy cows. For urine N excretion, body weight was the only significant factor beyond N intake in the present analysis, with a positive association between body weight and urine N excretion, which could reflect effects of total body protein turnover and 'non-productive' protein requirements. Yan et al. (2006) reported that body weight improved prediction of manure N excretion slightly when included with N intake in prediction models, but was a relatively poor predictor of manure N excretion on its own or in combination with milk yield. For lactating dairy cows only, both diet ash and ADF concentration were significantly associated with faecal N excretion. Diet ADF had a positive effect, perhaps also reflecting increased hindgut fibre fermentation or excretion of indigestible protein fractions. Diet ash concentration had a negative association with faecal and milk N excretion in lactating dairy cows, and also had a positive association with retained N. This suggests that diets with higher ash levels were associated with lower milk protein yield and thus relatively more body N retention, but the mechanisms involved are impossible to ascertain based on the diet composition data available in the current study. There was also a negative association between body weight and milk N excretion, as well as N retention in the dairy cow data base. Reasons for these associations between body weight and N excretion and retention, whereby larger cows excrete less dietary N increments in milk and retain less increased N intake in body tissues are not certain, but in a previous meta-analysis of N excretion and retention in growing beef cattle there was also a negative association between body weight and retained N as a proportion of N intake (Angelidis et al., 2019).

Initial visualisations of the data (figs 2a-e, 3a-e, 4a-e) suggested there to be variation associated with site and a number of methodologies associated with the measurement of nitrogen balance, however none of the methods identified were significant in the mixed models analysis and thus were not retained in the series of 'best' models produced for retained N, urine N or faecal N. A number of explanations exist for this apparent lack of relationship between methods and N excretion. For a number of the variables there was a dominant method in use and relatively few observations in the remaining category or categories such as, the measurement of DMI where manual weighing

accounted for more than 95% of observations and total collection of urine and faeces accounted for 92% of observations (table 1). In addition, these more uncommon methodologies were often only undertaken at a limited number of locations leading to confounding with site e.g. electronic feeding was only used in trials at 2 locations. Confounding with site was likely to have been a significant problem with the variable duration of collection, where many locations favour a single duration of collections and there is a lack of within site variation. The number of observations for 2, 3, 4 and 8 days collections are very small (table 1) so it is not possible to infer from this database if the variation observed at the extremes of collection periods is associated with variation in N measurement (figs 2e, 3e, 4e). There is relatively less variation associated with the more common collection durations of 5-7 days.

For all models except for faecal dry matter excretion in lactating dairy cows over half of the variation was associated with site and experiment. Experiment variation ranged from 28-37%, for retained N, 33-42% for urine N and was 26% for Milk N. Site variation ranged from 21-27%, for retained N, 20-25% for urine N and was 34% for Milk N. Site level variation was less for faecal N measurement (fig 4a) and this variation was interrelated with experiment variation, so it was not possible to run the models with experiment nested within experiment as a random effect. Experiment level variation was 56-62% for faecal N but the residual model variation was similar to that for the models of retained, urine and milk N. Similarly, site level excretion in faecal dry matter excretion was relatively low for both models, whilst experiment level variation in the lactating dairy cow model was also low. When compared with the site level variation for observed methane measurement of 7% obtained from a similar database meta-analysis, the site level variation associated with the measurement of N excretion in urine and milk and N balance is greater, reflecting the complexity of the data collection with multiple measurements begin taken, some manually, in order to calculate the final values of N intake, urine and milk N excretion, and finally retained N. For milk N excretion, the variation across sites likely reflects differences in milk yield of the cows used in the studies included.

For faecal dry matter excretion there was less variation due to site than observed for other components of N balance, suggesting that there is less variation associated with the methods used for collecting, weighing, sampling and analysing faecal dry matter. This in part reflects the limited number of steps involved and the relative stability of dry matter as a diet and faecal component. This does not account for variation diet or faecal volatile losses, which were not accounted for in the measurements of dry matter concentration evaluated in the current analysis.

For the methodology related variables, it is highly likely that there is a finer level of detail that was not captured adequately e.g. the N analysis method variable did have detail on how the faecal and urine samples were gathered, stored and prepared for analysis, and it is known that drying and grinding can result in volatile N loss from samples (Sphanghero and Kowalski, 1997; Morris et al., 2019). Also not captured was any data regarding the influence of human behaviour with regards to the methods used, which would be a significant challenge to collect but has the potential to have a considerable influence on N balance measurements. The ability to identify individual sources of variation that explain the high site level variation in N balance measurements was likely affected by lack of within site variation in methodologies, insufficient data for some methodologies, insufficient detail for some methodologies, and human behaviour in the conduct of the methods used. It is likely that a multi-site experimental protocol including a range of defined methodologies would be needed to successfully unpick the interactions between site and methodology.

Conclusions

As observed previously, dietary N intake is the major determinant of N excretion in faeces, urine and milk, but other dietary and animal factors modify the effects of N intake to varying degrees, ultimately reflecting effects of diet component intakes and nutrient supply relative to requirements. The elucidation of these factors was not the objective of the present meta-analysis, but the effects of diet and animal factors were accounted for in order to evaluate variation in measurements of N balance associated with methodology and research site. Considerable variation in measurements of N excretion and retention due to research site was observed, but this variation across sites was not due to the specific methodologies assessed and was not associated with similar variation in faecal dry matter and N excretion, which varied less due to site per se. The present analysis suggests that in addition to differences in the production of the experimental animals studied, variation across sites may be due more to how the specific experimental methods are used, rather than the method used per se. In this regard strict attention to detail in the conduct of methods and procedures and training of staff and students undertaking the experimental procedures at all levels, from feeding and collection of samples, through to sample storage, processing and analysis is critical (Snedecor and Flatt, 1970).

References

- Angelidis, A., Crompton, L., Misselbrook, T., Yan, T., Reynolds, C. K., Stergiadis, S. Evaluation and prediction of nitrogen use efficiency and outputs in faeces and urine in beef cattle. *Agriculture, Ecosystems and Environment*, 2019, 280, 1-15. doi.org/10.1016/j.agee.2019.04.013.
- Bougouin, A, A. Hristov, M.J. Aguerre, S. Ahvenjärvi, C. Arndt, A. Bannink, A.R. Bayat, C. Benchaar, T. Boland, W.E. Brown, L. A. Crompton, F. Dehareng, J. Dijkstra, I. Dufrasne, M. Eugène, E. Froidmont, S. van Gastelen, P.C. Garnsworthy, A. Halmemies-Beauchet-Filleau, S. Herremans, P. Huhtanen, M. Johansen, A. Kidane, M. Kreuzer, B. Kuhla, F. Lessire, P. Lund, E.M.K. Minnee, C. Muñoz, M. Niu, P. Nozière, D. Pacheco, E. Prestløkken, C. K. Reynolds, A. Schwarm, K. J. Shingfield, J.W. Spek, M. Terranova, A. Vanhatalo, M.A. Wattiaux, M.R. Weisbjerg, D. R. Yáñez-Ruiz, Z. Yu, and E. Kebreab. 2022. Prediction of nitrogen excretion from data on dairy cows fed a wide range of diets compiled in an intercontinental database: a meta-analysis. *J. Dairy Sci.*, accepted.
- Hristov, A. N., A. Bannink, L. A. Crompton, P. Huhtanen, M. Kreuzer, M. McGee, P. Nozière, C. K. Reynolds, A. R. Bayat, D. R. Yáñez-Ruiz, J. Dijkstra, E. Kebreab, A. Schwarm, K. J. Shingfield, Z. Yu. Nitrogen in ruminant nutrition: a review of measurement techniques. *Journal of Dairy Science*, 2019, 102, 5811-5852. 10.3168/jds.2018-15829
- Martin, A. K. 1966. Some errors in the determination of nitrogen retention of sheep by nitrogen balance studies. *Br. J. Nutr.* 20:325-337.
- Morris, D. L., A. W. Tebbe, W. P. Weiss, and C. Lee. 2019. Short communication: Effects of drying and analytical methods on nitrogen concentrations of feeds, feces, milk, and urine of dairy cows. *J. Dairy Sci.* 102:5212-5218.
- R Core Team (2020). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL
- Schneider, B. H., and W. P. Flatt. 1975. The Evaluation of Feeds through Digestibility Experiments. The University of Georgia Press, Athens, GA.
- Spanghero, M., and Z. M. Kowalski. 1997. Critical analysis of N balance experiments with lactating cows. *Livest. Prod. Sci.* 52:113-122.
- Spanghero, M., and Z. M. Kowalski. 2021. Updating analysis of nitrogen balance experiments in dairy cows. *J. of Dairy Sci.* Volume 104:7725-7737. <https://doi.org/10.3168/jds.2020-19656>.

van Lingen, H.J., Niu, M., Kebreab, E., Valadares Filho, S.C., Rooke, J.A., Duthie, C.-A., Schwarm, A., Kreuzer, M., Hynd, P.I., Caetano, M., Eugène, M., Martin, C., McGee, M., O’Kiely, P., Hünerberg, M., McAllister, T.A., Berchielli, T.T., Messana, J.D., Peiren, N., Chaves, A.V., Charmley, E., Cole, N.A., Hales, K.E., Lee, S.-S., Berndt, A., Reynolds, C.K., Crompton, L.A., Bayat, A.-R., Yáñez-Ruiz, D.R., Yu, Z., Bannink, A., Dijkstra, J., Casper, D.P., Hristov, A.N., 2019. Prediction of enteric methane production, yield and intensity of beef cattle using an intercontinental database. *Agric. Ecosyst. Environ.* 283, 106575. <https://doi.org/10.1016/j.agee.2019.106575>

Yan, T., J. P. Frost, R. E. Agnew, R. C. Binnie, and C. S. Mayne. 2006. Relationships among manure nitrogen output and dietary and animal factors in lactating dairy cows. *J. Dairy Sci.* 89:3981–3991.

Appendix 1. Correlation matrix for numerical variables ($R^2 > 0.5$ = yellow, $R^2 > 0.75$ = red)

	DMI	Forage %	CP cont	EE cont	ASH cont	NDF cont	ADF cont	STA cont	CP intake	N intake	EE intake	ASH intake	NDF intake	ADF intake	STA intake	DM dig	OM dig
DMI	1.00																
pcFor	0.00	1.00															
CP cont	0.22	-0.07	1.00														
EE cont	0.23	0.02	0.14	1.00													
ASH cont	0.00	0.29	0.23	-0.01	1.00												
NDF cont	-0.12	0.43	-0.17	-0.03	0.46	1.00											
ADF cont	0.07	0.48	-0.08	0.04	0.44	0.84	1.00										
STA cont	-0.41	-0.59	-0.21	-0.23	-0.43	-0.51	-0.65	1.00									
CP intake	0.93	0.01	0.54	0.25	0.07	-0.21	0.00	-0.43	1.00								
N intake	0.92	0.01	0.35	0.25	0.08	-0.20	0.00	-0.43	1.00	1.00							
EE intake	0.74	0.11	0.18	0.83	-0.03	-0.02	0.13	-0.42	0.72	0.72	1.00						
ASH intake	0.82	0.14	0.35	0.24	0.39	0.01	0.22	-0.50	0.84	0.84	0.68	1.00					
NDF intake	0.85	0.22	0.11	0.20	0.13	0.32	0.43	-0.56	0.76	0.76	0.64	0.76	1.00				
ADF intake	0.84	0.25	0.12	0.24	0.15	0.29	0.52	-0.62	0.75	0.75	0.67	0.76	0.93	1.00			
STA intake	0.65	-0.14	0.04	-0.01	-0.23	-0.36	-0.25	0.23	0.60	0.60	0.32	0.51	0.44	0.42	1.00		
DMdig	-0.10	-0.20	0.11	-0.01	-0.13	-0.38	-0.47	0.24	-0.03	-0.03	-0.08	-0.12	-0.27	-0.32	-0.16	1.00	
OMdig	-0.04	0.14	0.24	0.13	0.09	-0.12	-0.33	-0.15	0.08	0.08	0.00	0.01	-0.15	-0.30	-0.12	0.66	1.00
Ndig	0.02	-0.01	0.47	0.08	0.11	-0.11	-0.10	0.03	0.17	0.17	0.07	0.03	-0.08	-0.06	-0.04	0.37	0.53
NDFdig	-0.05	0.24	0.22	0.16	0.38	0.33	0.07	-0.31	-0.03	-0.03	0.00	0.07	0.05	-0.06	-0.38	0.31	0.69
ADFdig	0.17	0.33	0.14	0.22	0.32	0.26	0.24	-0.46	0.19	0.20	0.21	0.31	0.28	0.26	-0.14	0.15	0.63
STAdig	0.06	0.33	-0.03	-0.03	0.11	0.03	0.19	-0.23	0.03	0.03	0.11	0.09	0.06	0.09	-0.08	0.14	0.27
fecN excr	0.88	-0.03	0.22	0.15	0.06	-0.11	0.10	-0.37	0.85	0.85	0.62	0.78	0.78	0.79	0.57	-0.23	-0.29
urine vol	0.55	0.33	0.32	0.18	0.37	0.09	0.30	-0.55	0.58	0.58	0.48	0.60	0.52	0.56	0.31	-0.12	0.04
urineN excr	0.62	-0.04	0.23	0.18	0.13	-0.12	-0.01	-0.39	0.77	0.76	0.57	0.62	0.54	0.54	0.46	0.04	0.10
milk prodn	0.77	-0.32	0.21	0.18	-0.32	-0.49	-0.39	0.14	0.67	0.66	0.56	0.49	0.43	0.38	0.49	0.13	-0.01
milk fat	-0.24	0.27	0.10	-0.08	0.18	0.23	0.15	-0.18	-0.16	-0.15	-0.20	-0.10	-0.11	-0.17	-0.26	-0.03	0.12
milk lact	0.07	-0.07	-0.06	-0.10	-0.10	-0.11	-0.05	0.13	0.00	0.00	-0.11	-0.04	-0.02	-0.10	0.07	0.02	-0.01
MUN	-0.01	-0.14	0.14	0.10	0.13	-0.11	-0.21	0.08	0.06	0.06	0.06	0.07	-0.14	-0.13	0.03	0.08	0.08
milkN	0.79	-0.43	0.32	0.08	-0.36	-0.59	-0.45	0.16	0.71	0.68	0.53	0.47	0.42	0.49	0.54	0.13	-0.04
Nbal	0.11	0.02	0.08	0.18	0.07	0.02	0.08	-0.07	0.21	0.22	0.23	0.12	0.09	0.17	0.02	0.05	0.20
Faecal N%	-0.04	0.01	-0.25	-0.15	-0.06	0.17	0.20	-0.07	-0.20	-0.22	-0.12	-0.03	0.09	0.10	0.01	-0.40	-0.57
Urine N%	-0.44	-0.18	0.00	-0.22	0.12	-0.11	-0.18	0.29	-0.33	-0.33	-0.39	-0.34	-0.52	-0.52	-0.31	0.24	0.15
Manure N%	-0.45	-0.15	-0.12	-0.31	0.10	-0.01	-0.08	0.28	-0.43	-0.43	-0.47	-0.35	-0.47	-0.49	-0.33	0.05	-0.21
Milk N%	0.05	-0.05	-0.13	0.07	-0.42	-0.15	-0.17	0.10	-0.24	-0.25	0.07	-0.18	0.06	-0.02	0.12	0.02	-0.06
Retained N%	-0.14	-0.10	0.07	0.05	-0.04	-0.03	-0.03	0.18	-0.09	-0.04	-0.08	-0.16	-0.19	-0.14	-0.13	0.12	0.20
body weight	0.73	0.04	0.07	0.26	0.12	0.10	0.19	-0.47	0.64	0.63	0.61	0.63	0.69	0.68	0.40	-0.16	0.00

	N dig	NDF dig	ADF dig	STA dig	fecN excr	urine vol	urineN excr	milk prodn	milk fat	milk lact	MUN	milkN	Nbal	Faecal N%	Urine N%	Manure N%	Milk N%	Retained N%	body weight
Ndig	1.00																		
NDFdig	0.21	1.00																	
ADFDig	0.10	0.86	1.00																
STAdig	0.19	-0.06	-0.09	1.00															
fecN excr	-0.24	-0.14	0.13	-0.04	1.00														
urine vol	0.04	-0.06	0.18	0.21	0.49	1.00													
urineN excr	0.32	-0.11	0.01	0.03	0.54	0.64	1.00												
milk prodn	0.04	-0.19	-0.10	-0.08	0.58	0.18	0.33	1.00											
milk fat	0.02	0.28	0.28	0.07	-0.15	0.00	-0.10	-0.41	1.00										
milk lact	-0.08	-0.03	0.04	-0.03	0.03	0.06	-0.12	0.17	-0.09	1.00									
MUN	-0.02	0.05	-0.04	0.08	0.08	-0.17	0.06	0.01	-0.04	-0.13	1.00								
milkN	0.05	-0.37	-0.35	0.00	0.59	0.24	0.40	0.93	-0.36	-0.01	-0.09	1.00							
Nbal	0.30	0.14	0.18	0.00	-0.02	-0.04	-0.12	-0.07	-0.01	-0.01	0.04	-0.10	1.00						
Faecal N%	-0.90	-0.23	-0.08	-0.17	0.27	-0.09	-0.32	-0.07	0.00	0.05	0.01	-0.06	-0.45	1.00					
Urine N%	0.36	-0.05	-0.30	0.08	-0.43	-0.05	0.30	-0.12	0.03	-0.11	0.03	-0.01	-0.39	-0.25	1.00				
Manure N%	-0.11	-0.14	-0.30	0.01	-0.26	-0.12	0.10	-0.16	0.08	-0.08	0.04	-0.07	-0.63	0.31	0.84	1.00			
Milk N%	-0.28	-0.04	0.03	0.04	-0.06	-0.22	-0.38	0.45	-0.17	0.18	-0.08	0.37	-0.47	0.33	-0.31	-0.07	1.00		
Retained N%	0.27	0.15	0.11	-0.07	-0.23	-0.31	-0.28	-0.12	0.02	0.02	0.02	-0.14	0.87	-0.46	-0.33	-0.57	-0.50	1.00	
body weight	0.01	0.04	0.17	0.17	0.61	0.52	0.50	0.25	-0.10	-0.25	0.01	0.44	0.06	0.00	-0.26	-0.26	-0.09	-0.22	1.00